

THE DETECTION OF VARIABILITY FROM THE CANDIDATE IR COUNTERPART TO THE ANOMALOUS X-RAY PULSAR 1E 1048.1–5937¹

G.L. ISRAEL^{2,3}, S. COVINO⁴, L. STELLA^{2,3}, S. CAMPANA^{4,3}, G. MARCONI⁵, S. MEREGHETTI⁶, R. MIGNANI⁷, I. NEGUERUELA⁸, T. OOSTERBROEK⁹, A.N. PARMAR⁹, L. BURDERI²

AND

L. ANGELINI¹⁰
gianluca@mporzio.astro.it

Received 2002 August 12; accepted 2002 September 27

ABSTRACT

We report on the detection of variability from the proposed IR counterpart to the Anomalous X-ray Pulsar (AXP) 1E 1048.1–5937 based on Chandra and ESO optical/IR deep observations carried out in 2001–2002. Within the narrow *Chandra* uncertainty region for 1E 1048.1–5937 we found only one relatively faint ($J=22.1\pm0.3$, $J-K_s=2.4$) source, while the recently proposed IR counterpart was not detected down to a limiting K_s magnitude of ~ 20.7 (3σ confidence level). This implies a remarkable IR brightening of this object, $\Delta K_s > 1.3$, on a timescale of about 50 days. Although our knowledge of the IR properties of AXPs is rather limited (there is only another source, 1E 2259+586, for which IR variability has been detected), the observed IR variability of the proposed counterpart strengthens its association with 1E 1048.1–5937. Our results make the IR (and presumably optical) variability a likely common characteristic of AXPs, and provide new constraints on this class of objects.

Subject headings: stars: pulsars: general – pulsar: individual: – 1E 1048.1–5937 – infrared: stars – stars: variables: other – X-rays: stars

1. INTRODUCTION

After more than 20 years since the discovery of X-ray pulsations from the prototype source 1E 2259+586, the current observational properties of AXPs are not yet conclusive to unambiguously assess their nature. There are currently five confirmed members of the AXP class plus two candidates. Although we can be reasonably confident that AXPs are magnetic rotating neutron stars (NSs), their energy production mechanism is still uncertain. It is also unclear whether they are solitary objects (either low magnetised or magnetars) or are in binary systems with very low mass companions (for a review see Israel et al. 2002; Mereghetti et al 2002 and references therein). Different production mechanisms for the observed X-ray emission have been proposed, involving either accretion or the dissipation of magnetic energy. The recent detection of X-ray bursts from 1E 2259+586 and 1E 1048.1–5937 has strengthened the possible connection of AXPs with Soft Gamma-ray Repeaters (SGRs; Kaspi & Gavril 2002; Gavril et al. 2002). Moreover the detection of relatively large pulsed fraction pulsations in the optical flux of 4U 0142+614 (at the same period of X-ray pulses) seems to rule out the X-ray reprocessing and favors models based on isolated NSs (Kern & Martin 2002). The magnetar model, originally proposed by Thompson & Duncan (1993) to explain the properties of SGRs, appears currently to

be fairly successful at interpreting these properties of AXPs.

As done in the past in the case of High and Low Mass X-ray binary system (HMXBs and LMXBs), finding the optical/IR counterparts to AXPs might give the key to definitively assess the nature and the emission mechanisms of AXPs. The lack of any known distinctive property of AXPs in the optical/IR bands, makes the X-ray positional accuracy at the (sub)arcsec level a fundamental preliminary step to carry out very deep searches for potential optical/IR counterparts. Moreover a high X-ray spatial resolution can be exploited to search for small scale diffuse emission or the presence of structures in the vicinity of the sources.

1E 1048.1–5937 has been extensively studied in the X-ray band since its discovery in 1979 (Seward et al. 1986). Previous searches for an optical counterpart gave negative results (Seward et al. 1986; Mereghetti et al. 1992; Wang & Chakrabarty 2002a). Recently, a relatively faint IR counterpart ($J=21.7$; $J-K_s=2.3$) to 1E 1048.1–5937, with unusual IR colors, has been proposed (Wang & Chakrabarty 2002b; hereafter WC02).

In this paper we report on the early results obtained for 1E 1048.1–5937 as a part of a joint ESO/Chandra large project aimed at the identification and study of the optical/IR counterparts of AXPs in the southern sky, and the study of the spatial distribution of X-ray emission from AXPs (results concerning *Chandra* observations of 1RXS J170849–400910 and

¹ The results reported in this Letter are partially based on observations carried out at ESO, La Silla, Chile (66.D-0440 and 68.D-0350).

² INAF – Osservatorio Astronomico di Roma, V. Frascati 33, I-00040 Monteporzio Catone (Roma), Italy; gianluca, stella and burderi @mporzio.astro.it

³ Affiliated to the International Center for Relativistic Astrophysics

⁴ INAF – Osservatorio Astronomico di Brera, Via Bianchi 46, I-23807 Merate (Lc), Italy; covino and campana @merate.mi.astro.it

⁵ European Southern Observatory, Casilla 19001, Santiago, Chile; gmarconi@eso.org

⁶ Istituto di Fisica Cosmica G. Occhialini, CNR, Via Bassini 15, I-20133 Milano, Italy; sandro@mi.iasf.cnr.it

⁷ European Southern Observatory, Karl-Schwarzschildstr. 2, D-85748 Garching, Germany, rmignani@eso.org

⁸ Dpto. de Física, Ingeniería de Sistemas y Teoría de la Señal, Universidad de Alicante, Apdo. de Correos 99, E03080, Alicante, Spain; ignacio@astronomia.disc.ua.es

⁹ Astrophysics Missions Division, Research and Scientific Support Department of ESA, ESTEC, Postbus 299, NL-2200 AG Noordwijk, The Netherlands; toosterb and aparmar @rssd.esa.int

¹⁰ Laboratory of High Energy Astrophysics, Code 660.2, NASA/Goddard, Space Flight Center, MD 20771, USA; angelini@davide.gsfc.nasa.gov

AX J1844–0258 will be reported elsewhere).

2. *Chandra* OBSERVATIONS

The field including the X-ray position of 1E 1048.1–5937 was observed during *Chandra* Cycle 2. Previously determined positions of the source had limited accuracy; $15''$ for *EXOSAT* (Mereghetti et al. 1992) and $\sim 4''$ for *XMM-Newton* (Tiengo et al. 2002). The source was observed with the High Resolution Imager (HRC-I; Zombeck et al. 1995) in order to obtain the most accurate X-ray position. 1E 1048.1–5937 was observed with *Chandra* on 2001 July 4 for an effective exposure time of 9870 s. Data were reduced with CIAO version 2.2 and analysed with standard software packages for X-ray data (CIAO, XIMAGE, XRONOS, etc.). The observation was carried out with a nominal aspect solution and latest calibration files were used. Four different detection algorithms were used: the CIAO *wavdetect* and *celldetect* tool, the XIMAGE sliding cell, and the wavelet-based *pwdetect* software developed and optimised for *Chandra* images (Damiani et al. 1997). Based on a statistical study performed on several point-like sources detected within the field of view of public HRC-I archival data with known optical counterparts, we found a difference up to $0''.2$ in coordinates among the four different detection methods. In the following, we refer to the coordinates obtained with the latter method (we assumed that differences in the coordinates among the four methods are purely statistical).




FIG. 1.— Radial surface brightness profile of the *Chandra* HRC-I emission from a circular region of $7''$ radius around 1E 1048.1–5937. The dashed line represents the simulated *Chandra* point spread function for a point-like on-axis source.

Only one source was detected in the HRC-I. This has the following coordinates: R.A. = $10^{\text{h}}50^{\text{m}}07^{\text{s}}.12$, DEC. = $-59^{\circ}53'21''.37$ (equinox J2000), with an uncertainty circle radius of $0''.7$ (90% confidence level). Note that the *Chandra* position is not consistent with that obtained with *XMM-Newton* (Tiengo et al. 2001; R.A. = $10^{\text{h}}50^{\text{m}}06^{\text{s}}.3$, DEC. = $-59^{\circ}53'17''$; equinox J2000, 90% confidence level circle radius of $4''$). However, inspection of the attitude data during the *XMM-Newton* observation, showed that an error radius of about $9''$ is likely more realistic (Tiengo 2002, private communication). Photon

arrival times were extracted from a circular region centered on the previous reported position with a radius of $1''.5$, including more than 90% of the source photons. These were corrected to the barycenter of the solar system and timing analysis applied (CIAO *fxbary* tool). A coherent pulsation at a period of 6.4529 ± 0.0001 s was detected confirming that the detected source is indeed the AXP 1E 1048.1–5937. The latter value falls within the expected period interval extrapolated from the more accurate timing analysis obtained from a long-term *RXTE* monitoring program of 1E 1048.1–5937 (Kaspi et al. 2001; we assumed the maximum and minimum $\dot{P}$ reported in their Table 1).

One of the aims of the *Chandra* HRC-I observations of AXPs, was to study the radial profile of the X-ray emission from the AXPs with the unprecedented spatial resolution offered by the HRC-I. Figure 1 shows the spatial profile we obtained for the (background subtracted) X-ray emission from a region of $7''$ radius centered on the above reported source position. The radial profile is in good agreement with the expected *Chandra* point spread function (solid line simulated with MARX; Wise et al. 1997).

3. ESO OBSERVATIONS

Over the last two years we carried out deep optical and IR observations of the field including the original *Einstein* position of 1E 1048.1–5937. These data were obtained from the 3.5-m New Technology Telescope (NTT, La Silla, Chile) equipped with the Superb Seeing Imager-2 (SUSI2; 2 CCDs of 2048×4096 pixels; pixel scale of $0''.16$ and $5.5' \times 5.5'$ field of view) in the optical band, and with SOFI Isaac (SOFI; Hawaii HgCdTe 1024×1024 array; $0''.292$ pixel scale and $4.9' \times 4.9'$ field of view) in the near-IR band.

Optical Johnson filter deep images (BVRI) were first carried out with SUSI2 in service mode on 2001 March 20 with the following effective exposure times: 2700 s (B), 4350 s (V), 2400 s (R) and 3400 s (I). Weather conditions were good with seeing of about $1''$. Standard reduction packages were applied to the optical data (DAOPHOT II; Stetson 1987) to obtain the photometry of each stellar object in the images. Limiting magnitudes of 25.5 (B), 25.5 (V), 24.5 (R) and 23.5 (I) were reached.

JH images were initially carried out with SOFI in service mode on 2001 May 29; 2700 s (each filter) of effective exposure time. Unfortunately, during observations clouds were present and the seeing was poor ($1''$ – $1''.5$) yielding a limiting magnitude (3σ confidence level) of 22.5 (J) and 21 (H). SOFI JHKs images were obtained with better weather (photometric) and seeing ($0''.6$) conditions on 2002 February 19. Final exposure times were of 3600 s (J), 4200 s (H) and 4800 s (Ks). Single 10 s-long exposure images were taken for each filter (5 s-long in Ks) with offsets of $40''$ in order to sample and subtract the variable IR background. Standard IR software packages were used for sky frame subtraction and image coaddition (Eclipse and IRDR; Devillard 1997 and Sabbey et al. 2001). During the latter run limiting magnitudes of $J \sim 23$, $H \sim 21.5$, and $Ks \sim 20.7$ were reached.

To register the *Chandra* coordinates of 1E 1048.1–5937 on our optical/IR images, we computed the image astrometry using as reference the positions of stars selected from the USNO-A2.0 catalog which has an intrinsic absolute accuracy better than $0''.25$ (depending on magnitude and sky position of reference stars; Assafin et al. 2001). About 40 objects with mag-




FIG. 2.— Optical *I* band (left panel) and near-IR *Ks* band (right panel) images of the region including the position of 1E 1048.1–5937, taken from the 3.5-m ESO NTT. The *Chandra* uncertainty circles are also shown: the thick solid line circle has radius $0''.8$ and represents the position reported in this paper, while the thin one is the WC02 position with a radius of $1''$. The faint object marked with X3, which lies on the upper edge of the circles, is the only one consistent with the X-ray position. The square represents the position expected of X1, proposed as the IR counterpart to 1E 1048.1–5937 by WC02. Coordinates are RA (h m s) and Dec. ($^{\circ}$ ' ' ' ; equinox J2000).

nitudes brighter than $R=14$ were identified in the optical and IR images and used for astrometric calibrators. The pixel coordinates of the reference stars were computed by a two dimensional Gaussian fitting procedure and transformation from pixel to sky coordinates was then computed using the Graphical Astronomy and Image Analysis Tool (GAIA,¹¹). After taking into account the uncertainties of the source X-ray coordinates ($0''.7$), the rms error of our astrometry ($0''.15$) and the propagation of the intrinsic absolute uncertainties on the USNO–A2.0 coordinates ($0''.25$), we estimated the final accuracy to be attached to the 1E 1048.1–5937 position as about $0''.8$. Figure 2 shows a region of $20'' \times 20''$ around the 1E 1048.1–5937 position in the *I* and *Ks* bands with the *Chandra* uncertainty circle superimposed (90% confidence level).

4. DISCUSSION

The NTT IR data show that no object is consistent with the *Chandra* position of 1E 1048.1–5937 in the optical band down to a *V* limiting magnitude of about 25.5 (see Figure 2). On the other hand, a faint object (marked X3 in Figure 2) is present in the IR *Ks* frames ($J=22.1 \pm 0.3$, $Ks=19.7 \pm 0.3$). This object is also detected in the WC02 frames, although its magnitudes are not listed. A visual inspection of the WC02 image suggests that the source X3 has remained nearly unchanged in flux with respect to our frames (superimposing to our IR images the *Chandra* position and uncertainty radius reported by WC02 for 1E 1048.1–5937, the object X3 falls within the circle). Note that during the NTT IR observations carried out on 2001, the object X3 is not detected due to the poor seeing conditions and contamination with the nearby bright object. More surprisingly, we note that no object is detected at the position of the source X1 reported by WC02 (the position of X1 is marked with a square in Figure 2). For comparison, the object marked X2 in

Figure 2 has a magnitude of $J=20.63 \pm 0.06$, $H=19.73 \pm 0.15$ and $Ks=19.0 \pm 0.2$ (consistent, within $1-1.5\sigma$, with the values reported by WC02). As the limiting magnitudes of our IR frames are well above the values measured by WC02 for X1, we conclude that the source was considerably fainter during our observations. This IR brightening ($\Delta Ks > 1.3$; about a factor of 3 in flux), displayed by the source on a timescale of about 50 days (the time interval between our last IR images and those of WC02) further strengthens the link between the IR source and 1E 1048.1–5937. Moreover, the chance probability of finding a highly variable IR source within a circle of $0''.8$ is negligible.

An IR counterpart was recently found for the prototype of the AXP class, 1E 2259+586, which also showed IR variability at a level of $\Delta Ks \sim 1.3$ (Kaspi et al. 2002). This IR variability might be associated to the short-term X-ray bursts (similar to those displayed by SGRs) observed from the source three days before the IR observations (Kaspi & Gavril 2002). Recently, short X-ray bursts were detected in the X-ray flux of 1E 1048.1–5937 (Gavril et al. 2002). All these findings together, suggest a close similarity of 1E 1048.1–5937 and 1E 2259+586, such that the IR counterpart of the former might be expected to be variable. For this reason X1 is the most likely IR counterpart to 1E 1048.1–5937, even though source X3 is included in the *Chandra* uncertainty circle and its color ($J-Ks=2.4$) is within the value interval predicted for AXPs (see also Figure 2 of WC02).

The fact that 1E 1048.1–5937 showed two X-ray bursts close to each other, implies that many other bursts might have been missed due to the sparse X-ray monitoring with the *RXTE*. Moreover since our non-detection of the IR candidate predates the WC02 detection, it would appear more natural to associate (if at all) the IR variability to some X-ray activity that took place between the two IR campaigns (as opposed to the

¹¹ See <http://star-www.dur.ac.uk/~pdraper/gaia/gaia.html>

few second-duration X-ray bursts that was detected 4 months earlier). Finally we note that the X-ray/IR observations of 1E 2259+586 are such that we do not know whether the detected IR brightening follows or precedes that in the X-ray band. Future multi-wavelength monitoring programs will probably allow us to better understand these points.

The non-detection of X1 in our IR frames implies that the “quiescent” (or low state) IR emission level of 1E 1048.1–5937 is still undetected. Correspondingly, it cannot be ruled out yet that the spectral flattening proposed by WC02 might result from wavelength-dependent variability. Similarly, the measured IR fluxes for the counterpart cannot be used to study the overall (from X-ray to IR) energy spectrum of 1E 1048.1–5937, since they reflect a different energy distribution than the quiescent (or low state) one. Deeper IR observations are needed to determine the quiescent emission state of this source.

IR variability might represent a new, fairly common, characteristic of AXPs (and possibly SGRs), and potentially a diagnostic of their nature. We note that, so far, very little is known on the expected optical/IR emission from a magnetar. Both models based on magnetar and accretion from “fossil” disk have difficulties in accounting for transient and/or variable emission at wavelengths shorter than those in the X-ray band (Thompson & Duncan 1996 and references therein; Chatterjee & Hernquist 2000 and references therein). On the other hand, the analogy with the X-ray bursts of LMXBs, suggests that the

effects of the X-ray burst of AXPs might propagate towards longer wavelengths (e.g., via reprocessing). It is somewhat intriguing that at least the values of ΔK s and colors of the IR counterpart of 1E 2259+586 and 1E 1048.1–5937 are similar to those displayed by some classes of accreting cataclysmic variables and low mass X-ray binaries hosting a neutron star.

The superb *Chandra* HRC-I spatial resolution allowed us to look for extended X-ray emission around the pulsar; this might result from an X-ray nebula powered by the AXPs as well as a scattering halo. A first detailed study was carried out for 1E 2259+586, where emission beyond 4'' and extending up to 100'' was found (Patel et al 2001). However in this case the source is embedded in the diffuse emission from the supernova remnant G109.1–1.0, and it is very difficult to disentangle this component from that (possibly present) around the AXP. The present *Chandra* HRC-I data are by far the best spatial resolution observation carried out for an AXP and no significant diffuse X-ray emission was found beyond $\sim 2''$.

This work is supported through CNAA, ASI, CNR and Ministero dell'Università e Ricerca Scientifica e Tecnologica (MURST-COFIN) grants. The authors thank Olivier Hainaut, Leonardo Vanzì and the NTT Team for their kind help during ESO observations. We thank Fotis Gavriil for a careful reading of an early version of the paper.

REFERENCES

- Assafin, M., Andrei, A.H., Martins, R.V., da Silva Neto, D.N., Camargo, J.I.B., Teixeira, R., & Benevides-Soares, P. 2001, *ApJ*, 552, 380
 Chatterjee, P. & Hernquist, L. 2000, *ApJ*, 543, 368
 Damiani, F., Maggio, A., Micela, G., & Sciortino, S. 1997, *ApJ*, 483, 350
 Devillard, N. 1997, *The Messenger*, 87
 Gavriil, F.P., Kaspi, V.M., Woods, P.M. 2002, *Nature*, 419, 142
 Kaspi, V.M., Gavriil, F.P., Chakrabarty, D., Lackey, J.R., & Muno, M.P. 2001, *ApJ*, 558, 253
 Kaspi, V.M., Gavriil, F.P., & Woods, P.M. 2002a, *IAU Circ.*, 7926
 Kaspi, V.M. & Gavriil, F.P. 2002b, *IAU Circ.*, No. 7924
 Kern, B. & Martin, C. 2002, *Nature*, 417, 527
 Israel, G.L., Mereghetti, S., & Stella, L. 2002, in γ -Ray Bursts in the Afterglow Era, ed. S. Mereghetti & M. Feroci, *Mem.S.A.It.*, in press (see also astro-ph/0111093)
 Mereghetti, S., Caraveo, P., & Bignami, G.F. 1992, *A&A*, 263, 172
 Mereghetti, S., Chiarlone, L., Israel, G. L., & Stella, L. 2002, in *Neutron Stars, Pulsars and Supernova Remnants*, eds. W. Becker, H. Lesch & J. Trümper (Garching: MPE), in press (astro-ph/0205122)
 Patel, S.K. et al. 2001, *ApJ*, 563, L45
 Sabbey, C.N., McMahon, R.G., Lewis, J.R., & Irwin, M.J. 2001, *ASP Conf. Ser.* 238: *Astronomical Data Analysis Software and Systems X*, 10, 317
 Seward, F. D., Charles, P. A., & Smale, A. P. 1986, *ApJ*, 305, 814
 Stetson, P.B., 1987, *PASP*, 99, 191
 Thompson, C., & Duncan, R.C. 1993, *ApJ*, 408, 194
 Thompson, C., & Duncan, R.C. 1996, *ApJ*, 473, 322
 Tiengo, A., Göhler, E., Staubert, R., & Mereghetti, S. 2002, *A&A*, 383, 182
 Wang, Z.-X. & Chakrabarty, D. 2002a, *ASP Conf. Ser.* 271: *Neutron Stars in Supernova Remnants*, 297
 Wang, Z.-X. & Chakrabarty, D. 2002b, *ApJ*, in press (WC02)
 Wise, M.W., Huenemoerder, D.P., & Davis, J.E. 1997, *ASP Conf. Ser.* 125: *Astronomical Data Analysis Software and Systems VI*, 6, 477
 Zombeck, M.V., Chappell, J.H., Kenter, A.T., Moore, R.W., Murray, S.S., Fraser, G.W., & Serio, S. 1995, *Proc. SPIE*, 2518, 96